

Northeastern Interior Dry-Mesic Oak Forest



Macrogroup: Central Oak-Pine

This map is a modeled distribution based on current data and is not a substitute for field based inventory. Contact your State Natural Heritage Ecologist for more information about this habitat.



© Gary P. Fleming (Virginia Department of Conservation & Recreation Natural Heritage Program)

Description:

An oak-dominated, mostly closed canopy forest that occurs as a matrix (dominant) type through the central part of our region. Oak species characteristic of dry to mesic conditions (e.g., red, white, black, and scarlet oak) and hickories are dominant in mature stands. Chestnut oak may be present but is generally less important than other oak species. Red maple, black birch, and yellow birch may be common associates. Heath shrubs are often present but not well developed. Local areas of limy bedrock, or colluvial pockets, may support forests that reflect the richer soils. With a long history of human habitation, many of the forests are mid-successional, in which pines (typically Virginia or white) or tuliptree may be codominant or dominant.

State Distribution: CT, DC, DE, MA, MD, NJ, NY, PA, RI, VA, WV

Total Habitat Acreage: 17,032,701

Percent Conserved: 19.1%

State	State Habitat %	State Acreage	GAP 1&2 (acres)	GAP 3 (acres)	Unsecured (acres)
PA	37%	6,264,459	220,896	1,188,152	4,855,411
WV	22%	3,732,111	40,981	289,214	3,401,916
VA	15%	2,588,383	299,870	452,215	1,836,298
NY	11%	1,811,589	19,982	155,854	1,635,753
CT	6%	965,419	38,892	123,495	803,032
MD	4%	678,802	60,757	111,810	506,235
NJ	3%	559,819	117,260	47,837	394,722
MA	1%	242,876	5,771	34,365	202,741
RI	1%	179,468	8,231	29,188	142,049
DE	0%	8,229	59	2,573	5,596
DC	0%	1,546	0	0	1,546

Ecological Setting and Natural Processes:

Moderate moisture and heat loading are characteristic for this oaky system. It occurs at low to mid elevations, where the topography is flat to gently rolling, occasionally steep. Substrate bedrock and soils are commonly but not always acidic. Chestnut was formerly a prominent tree in these forests.

Similar Habitat Types:

Drier oak-pine systems (Central Appalachian Dry Oak-Pine Forest, CA Pine-Oak Rocky Woodland) are often upslope; mesic covey or wetland systems may be embedded in low landscape positions. A split along purely geographic lines separates this system from similar Southern Appalachian Oak Forests in southern WV, in lieu of more natural ecological or floristic distinctions.

Crosswalk to State Wildlife Action Plans:

Hardwood Forest - Mixed oak-beech forests (DC), Mesic Deciduous Forests (MD), Upland forests - deciduous forest (NJ), Oak Forest (NY), Deciduous/Mixed Forest (upland) (PA), Forest Habitat - Deciduous Forest (VA), Oak/Hickory and Dry/Mesic Oak Forest (WV)

Crosswalk to State Name Examples:

Dry Subacidic Forest (CT), Central Appalachian Dry-Mesic Chestnut Oak-Northern Red Oak Forest (DE), Dry, Rich Acidic Oak Forest (MA), Acidic Oak - Hickory Forest (MD), Dry-Mesic Inland Mixed Oak Forest (NJ), Appalachian Oak-Hickory Forest (NY), Dry Oak-Heath Forest (PA), Black Oak-Scarlet Oak/Heath Forest (RI), Central Appalachian Dry-Mesic Chestnut Oak - Northern Red Oak Forest (VA), Oak/Hickory And Dry/Mesic Oak Forest (WV)

Places to Visit this Habitat:

Green Ridge State Forest | MD
 Delaware Water Gap | NJ
 Sprout State Forest | PA
 George Washington and Jefferson National Forest | VA
 Monongahela National Forest | WV

Associated Species: *Appendix lists scientific names*

BIRDS: black-and-white warbler, broad-winged hawk, cerulean warbler, eastern wood-pewee, great crested flycatcher, louisiana waterthrush, ovenbird, red-bellied woodpecker, scarlet tanager, summer tanager (south), eastern whip-poor-will, wood thrush, veery, worm-eating warbler

MAMMALS: black bear, red-backed vole, short-tailed shrew, white footed mouse

HERPTILES: northern redback salamander, ringneck snake, redbelly snake, spotted salamander

PLANTS: American wintergreen (*Pyrola americana*), basil beebalm (*Monarda clinopodia*), blunt-lobe woodsia (*Woodsia obtusa*), bottlebrush grass (*Elymus hystrix*), common alexanders (*Zizia aurea*), early buttercup (*Ranunculus fascicularis*), shinleaf (*Pyrola elliptica*), sicklepod (*Arabis canadensis*)

Species of Concern (G1-G4): *Appendix lists scientific names*

BIRDS: golden-winged warbler

MAMMALS: eastern small-footed myotis, kittatiny red-backed vole, virginia big-eared bat


HERPTILES: big levels salamander, milk snake, peaks of otter salamander

INSECTS: American snout, Appalachian grizzled skipper, underwing moth (*Catocala resecta*), clouded underwing, dark stoneroot borer moth, flypoison borer moth, habilis underwing, northern metalmark, mournful underwing, yellow stoneroot borer moth

PLANTS: climbing fern (*Lygodium palmatum*), goldenseal (*Hydrastis canadensis*), small whorled pogonia (*Isotria medeoloides*)




© Gary P. Fleming (Virginia Department of Conservation & Recreation Natural Heritage Program)




The average patch size for this habitat is 13 acres and the largest single patch is 20,946 acres. This chart shows the proportion of the habitat that is in each patch-size class.



This chart shows the average age of trees associated with this habitat based on forest inventory data. For non-forested systems or small habitats the average age is influenced by the surroundings.



This chart shows the predicted loss of habitat over the next five decades (783,733 acres) if loss continues at the same rate as 1990-2000. The average rate of loss is 15,675 acres per year.



This metric measures how connected or fragmented the land directly surrounding (18 square miles) the habitat is, this the chart shows the proportion of the habitat in each connectedness class.